

Antibacterial Activity of Ethanol Extract, *n*-Hexane and Ethyl Acetate Fraction of Mundar (*Garcinia forbesii*) Pericarp

Hayatun Izma; Muhammad Ikhwan Rizki; Khoerul Anwar; Dyah Anggraeni; Satrio Wibowo Rahmatullah; Aditya Maulana Perdana Putra; Dita Ayulia Dwi Sandi

Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Universitas Lambung Mangkurat, Indonesia

ABSTRACT

Garcinia forbesii is a typical fruit of South Kalimantan, used by the community as medicine and cosmetics. The aimed of the study was to determine the antibacterial activity of ethanolic extract and fraction of *Garcinia forbesii* pericarp against *Staphylococcus aureus* and *Propionibacterium acne*. Antibacterial activity based on the diameter of inhibition zone, the minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC). The results showed that the largest inhibition zone diameter against *Staphylococcus aureus* and *Propionibacterium acne* was indicated by ethyl acetate fraction of 5.08 ± 1.020 mm and 14.33 ± 3.326 mm. The MIC value of ethanol extract against *Staphylococcus aureus* and *Propionibacterium acne* bacteria was $>1.5\%$, *n*-hexane and ethyl acetate fraction had the same results, namely against *Staphylococcus aureus* $>1.5\%$ and *Propionibacterium acne* at 1.5% . The MBC value was only obtained for fraction samples of *Propionibacterium acne* at concentration of 1.5% . It was concluded that ethyl acetate had strong antibacterial activity against *Propionibacterium acne* based on the value of inhibition zone diameter, MIC, and MBC.

Keywords: *Garcinia forbesii*, *Propionibacterium acne*, *Staphylococcus aureus*

INTRODUCTION

Bacteria are a group of single-celled living organisms and are microscopic (Prasetyo and Sasongko, 2014). *Staphylococcus aureus* and *Propionibacterium acne* are gram-positive bacteria (Rosyad, 2009; Karlina *et al.*, 2013; Prasetyo and Sasongko, 2014). *Staphylococcus aureus* causes pneumonia, lung abscess, meningitis, laryngeal infections, and skin lesions (Rostinawati, 2010; Amalia *et al.*, 2014). *Propionibacterium acne* causes pimples and plays a role in the inflammatory chemotactic process in the ducts of the sebaceous glands (Sambou *et al.*, 2017).

Antibacterials are compounds that can inhibit the growth of certain bacteria (Rostinawati, 2010), while antibiotics are used to treat bacterial infections (Amalia *et al.*, 2014). Antibiotic resistance cause problems that can counteract antibiotic treatment (Noviardi, 2010). The World Health Organization states that more than 200,000 newborns die yearly from infections that do not respond to the given drugs; most of these deaths occur in developing countries (Organization, 2021). Control of antibiotics resistance can be done by using plants that have antibacterial properties. One of the plants with the potential as an antibacterial agent is the pericarp of mundar (*Garcinia forbesii*).

The mundar (*Garcinia forbesii*) plant is a typical plant of South Kalimantan, traditionally used by the community as medicine and cosmetics (Alen *et al.*, 2008; Noor and Ningsih, 2017). The *Garcinia* genus contains secondary metabolite compounds such as xanthenes, alkaloids, flavonoids, tannins, and saponins (Ritthiwigrom *et al.*, 2013; Aziz, 2015). Alpha-mangostin is a compound belonging to the xanthone group found in the rind of the mangosteen fruit (*Garcinia mangostana* L.), and has been shown to have antibacterial activity (Sriyono and Andriani, 2013; Karthiga *et al.*, 2012; Miladiyah and Rachmawaty, 2017). This study aimed to determine the antibacterial activity of ethanol extract and fraction from *Garcinia forbesii* pericarp against *Staphylococcus aureus* and *Propionibacterium acne* based on the diameter of the

***Corresponding Author:** Hayatun Izma

Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Universitas Lambung Mangkurat

Email: hayatun.izma@ulm.ac.id

inhibition zone and determine the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC). The results of this study will increase the variety of plants that are scientifically proven to have antibacterial activity, and the results of this study also have the potential to be developed into new antibiotics derived from natural ingredients.

MATERIALS AND METHOD

Materials

The tools used were glassware (Pyrex® Iwaki Glass), measuring instruments (Pyrex® Iwaki Glass), autoclave (TOMY SX-500), stir bar, maceration vessel, blender (Miyako), bunsen, porcelain, separatory funnel, hot plate (MaxBlend), incubator (Mettler), *laminar air flow*, refrigerator, micropipette, analytical balance (Ohaus).

The ingredients used were *Garcinia forbesii* pericarp, distilled water, aluminum foil, bacteriological agar no. 1 (OXOID), barium chloride, ferric chloride, blank paperdisc, cotton swab, Chloramphenicol Supplement (OXOID), liquid spirits, 70% ethanol (technical), ethyl acetate (technical), gelatin, Whatman No.1 filter paper, bacterial culture *Staphylococcus aureus* ACC 25923, *Propionibacterium acne* ATCC 6919.

Methods

1. Plant Determination

Plant determination aims to match the morphological characteristics of the plants to be studied by looking at the literature, so there is no mistake in taking plants for research. Plant determination was carried out at the Basic Biology Laboratory, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, and Dr. Gunawan, S.Si., M.Si. as a botanist. Based on the test result certificate number 040a/LB.LABDASAR/I/2019 the results of the determination stated that the sample used in this study was *Garcinia forbesii*.

2. Sample Preparation

Samples of *Garcinia forbesii* pericarp were wet sorted, washed, and the pericarp was separated from the fruit flesh. Then cut into thin slices and dried at 60°C for three days. Dried samples were powdered using a grinder, then weighed and stored in a container. As much as 55 grams of dry powder of mundar pericarp was macerated with 70% ethanol at a ratio of 1:10 for 3 x 24 hours. The maceration results were filtered, the solvent was evaporated with a rotary evaporator, and concentrated with a waterbath (Anindya, 2012).

Fractionation using *n*-hexane and ethyl acetate using the liquid-liquid extraction. The extract was dissolved in aquadest, added with *n*-hexane, and leave it until two layers were formed. The *n*-hexane fraction was taken and concentrated using a waterbath. The aquadest fraction was added with ethyl acetate solvent, shaken, and left until two layers were separated. The ethyl acetate fraction was separated and then concentrated using a waterbath (Firdausi *et al*, 2015).

3. Test Sample Preparation

The samples used at concentrations of 1.5%, 1.25%, 1%, 0.75%, and 0.5% (b/v) from ethanol extracts, *n*-hexane fractions, and ethyl acetate fractions of *Garcinia forbesii* pericarp. The positive control used 1% chloramphenicol, while the negative control used DMSO because it was a diluent (Kaharap *et al*, 2016).

4. Bacterial Rejuvenation

The test bacteria *Staphylococcus aureus* ATCC 25923 and *Propionibacterium acne* ATCC 6919 came from the Banjarbaru Industrial Research and Standardization Center Laboratory. Each bacterium was

taken 1 mL and then inoculated into the Nutrient Broth (NB) medium in a closed test tube and incubated at 37°C for 24 hours (Poeloengan and Praptiwi, 2010). The bacteria were incubated then visually compared for turbidity with a 0.5 McFarland standard (10⁸ colonies/mL). Other comparisons were made by plating bacteria on NA media using the pour plate method. The bacteria were diluted, then plated, and put into a petri dish, and sufficient NA media was added and then incubated for 24 hours. The incubation results were observed, and counted manually how many bacteria grew (Weinstein, 2018).

5. Inhibition Zone Diameter Test

Inhibition zone by the diffusion method makes use of a paper disc or the so-called Kirby Bauer method. Nutrient agar medium is prepared by mixing 15 g of agar with 13 g of *nutrient broth* and adding distilled water to a volume of 1000 mL. Nutrient agar sterile medium is poured into the petri dish until evenly distributed and allowed to solidify. The petri dish containing the solidified media was divided into six areas, then the bacterial suspension was smeared on the Nutrient agar medium using a sterile *cotton swab*. A *blank paper disc* is dipped into each extract and fraction according to the concentration, then affixed to the media that has been bacteria smeared, and incubated for 24 hours at 37°C. The diameter of the clear zone formed was measured with a ruler and recorded (Rachmawaty, 2016).

The formula can calculate the diameter of inhibition zone: $\frac{d1+d2}{2} - x$.

Description:

d1 = vertical diameter of the clear zone in the media.

d2 = horizontal diameter of the clear zone in the media.

X = disc paper diameter (6 mm).

6. Minimum Inhibitory Concentration Test (MIC)

The MIC test was carried out using the microdilution method. The bacteria to be used are equated with the McFarland standard 0.5 (10⁵ colonies/mL), which is prepared by taking 1 mL of bacteria from the 10⁸ colonies/mL previously made and put into a test tube containing 9 mL of *nutrient broth*, then diluted again by taking 0.5 mL of bacteria and putting it into 9.5 mL of *nutrient broth*, so that 10⁵ colonies/mL of bacteria are obtained. 0.1 µL bacterial suspension was put into a *polypropylene PP centrifuge tube* containing 0.7 µL NB, then added with 0.2 µL sample (extract or fraction). This treatment was carried out on all *tubes*, the only difference being the concentration used (Weinstein, 2018), and then incubated for 24 hours at 37°C. After incubating, the *tube* containing the bacteria and the ethanol extract sample, the *n*-hexane fraction and the ethyl acetate fraction were smeared on the surface of the NA media in a petri dish divided into five parts, then incubated for 24 hours at 37°C. The minimum inhibitory concentration results are taken from the lowest concentration where there is no bacterial growth (Chikezie, 2017).

7. Minimum Bactericidal Concentration Test (MBC)

The MBC test was carried out using the *Streak plate* (scratch method). MBC determinations were carried out at concentrations considered as MIC values. The NA media was put into a petri dish and allowed to solidify. Samples with MIC values were taken to be scratched on the surface of the NA media using a *cotton swab*, then incubated for 24 hours at 37°C. Results that show no more bacterial growth are considered MBC. A clear area on the petri dish indicates no bacterial growth. In contrast, an area that looks like streaks indicates bacterial growth, so at this concentration, it can only inhibit bacterial growth (Rachmawaty, 2016; Chikezie, 2017).

DATA ANALYSIS

The effects of the inhibition zone diameter on bacterial growth inhibition of ethanol extract samples, *n*-hexane fractions, and ethyl acetate fractions were analyzed using SPSS (Wahyuddin, Pakadang and Aprilyani, 2017). The data in the second stage were in the form of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC). Each data is presented in tabular form and interpreted according to the results obtained.

RESULT AND DISCUSSION

1. Plant Determination Test Results

The results of the determination test showed that the sample used in this study was Mundar fruit (*Garcinia forbesii*) based on the test results certificate Number 040a/LB.LABDASAR/I/2019. The results are that the tree has a height of 15-17 m, leaves are oblong with a width of 6.63 cm, and stem height is 3-4 m. The fruit is spherical or round with a diameter of 2-3 cm, weighing less than 90 g; the color of the fruit is red when ripe, and the color of the fruit coating is a white cream with a soft and watery texture. The seeds are light brown with an oblong shape, 1.28 cm long, 0.65 cm wide, and 0.25 cm thick.

2. Sample Preparation Results

The dry powder result obtained was 43.39 g from the 500 g wet sample. These results indicate that there is shrinkage of the sample after drying. This is due to the large amount of water in the *Garcinia forbesii* pericarp. Previous research stated that the water content contained in *Garcinia forbesii* pericarp was 11% (Mranani, 2015).

Garcinia forbesii pericarp dry powder was extracted and then calculated the extract obtained. The yield calculation results are provided in table 1. Based on the table 1, the percent yield obtained in this study is not much different from previous studies. The percent yield of *Garcinia forbesii* pericarp obtained in previous research extracted using 70% ethanol had a yield of 57.83% (Dewi, 2018; Marsella, 2018). Calculation of percent yield was carried out to determine the percentage of metabolites extracted from the extraction process (Rizki *et al.*, 2021).

Table 1. Extraction results of *Garcinia forbesii* pericarp

Sample	Powder Weight	Extract Weight	Yield Value (%)
<i>Garcinia forbesii</i> ethanol extract	55.00 g	31.573 g	57.40

The extract obtained was then carried out by liquid-liquid fractionation. The results of *Garcinia forbesii* pericarp fractionation can be seen in Table 2. Based on the results of yield calculations, the ethyl acetate fraction has a more significant yield percentage than the *n*-hexane fraction. The yield calculation results are related to the number of metabolite compounds extracted in a sample. The sample's high content of metabolite compounds is indicated by the resulting high yield value (Sayuti, 2017). The percentage yield obtained from the *n*-hexane fraction of *Garcinia forbesii* pericarp was not much different when compared to the percent yield of the *n*-hexane fraction of mangosteen pericarp (*Garcinia mangostana*) which was 1.34% (Asifa, 2014), while the percent yield of the ethyl acetate fraction more significant than that of previous research, namely 17.08% in the ethyl acetate fraction of mangosteen pericarp (*Garcinia mangostana*) (Saputro, 2014; Asifa, 2014; Sayuti, 2017; Marsella, 2018; Rizki *et al.*, 2021). The results obtained differed from the percent yield of the ethyl acetate fraction of *Garcinia forbesii* pericarp in previous studies, which was 28.5% (Marsella, 2018).

Table 2. *Garcinia forbesii* pericarp fractionation results

Extract Weight	<i>n</i> -Hexane Fraction Weight	Ethyl Acetate Fraction Weight	<i>n</i> -Hexane Yield Fraction (%)	Ethyl acetate Yield Fraction (%)
20.00 g	0.201 g	14.01 g	1.005	70.05

3. Inhibitory Power Test Results

The results of the diameter of the inhibition zone of the samples of *Garcinia forbesii* pericarp can be seen in Table 3. The results showed that the diameter of the inhibition zone formed in the ethyl acetate fraction was more significant than in the ethanol extract and *n*-hexane fraction; this was probably due to the compounds contained in the ethyl acetate fraction of *Garcinia forbesii* pericarp being more active than those in the ethanol extract and the *n*-hexane fractions, and there were specific compounds act as antibacterial in the ethyl acetate fraction (Salni *et al*, 2011; Lestari *et al*, 2016). The samples of *Garcinia forbesii* pericarp had different inhibition zone diameters against *Staphylococcus aureus* and *Propionibacterium acne* bacteria.

Table 3. Diameter of zone inhibition in bacterial growth from samples of *Garcinia forbesii* pericarp

Sample	$x \pm SD$ diameter of zone inhibition (mm)	
	<i>S.aureus</i>	<i>P.acne</i>
Ethanol Extract	2.16±0.983	2.75±1.405
<i>n</i> -Hexane Fraction	2.08±0.664	4.08±0.664
Ethyl Acetate Fraction	5.08±1.020	14.33±3.326
Positive Control	6.24±0.592	19.5±1.449
Negative Control	0±0	0±0

Antibacterial activity can be influenced by several factors, including the concentration of the sample (extract or fraction), the content of antibacterial compounds, the diffusing power of the extract or fraction, the type of bacteria that is inhibited, the thickness of the agar plate, and the sensitivity of the bacteria to samples of *Garcinia forbesii* pericarp (Prihandani *et al.*, 2015). Based on the diameter of the inhibition zone, the positive control of 1% chloramphenicol proved that the antibiotic compound could affect positively to the growth of bacteria, while the negative control DMSO did not affect the growth of bacteria by not forming an inhibition zone.

The results showed that the activity as antibacterial of samples against *Staphylococcus aureus* bacteria was classified as weak. Based on the results of the Shapiro-Wilk test, the value of $p > 0.05$, or the data is normally distributed, the test is continued with the homogeneity test, the result is $p < 0.05$, or the data is not homogeneous. The analysis continued with a non-parametric test, namely Kruskal-Wallis and the Mann-Whitney test. The results of the Mann-Whitney test of the samples on the growth of *Staphylococcus aureus* bacteria can be seen in Table 4.

Table 4. Mann-Whitney test results for the samples of *Garcinia forbesii* pericarp against *Staphylococcus aureus* bacteria based on the diameter of the inhibition zone

Treatment group	Aquadest	Ethanol extract	<i>n</i> -hexane fraction	Ethyl acetate fraction	Chloramphenicol 1%
Aquadest	-	0.002*	0.002*	0.002*	0.002*
Ethanol extract	0.002*	-	0.738	0.004*	0.004*
<i>n</i> -hexane fraction	0.002*	0.738	-	0.004*	0.004*
Ethyl acetate fraction	0.002*	0.004*	0.004*	-	0.053
Chloramphenicol 1%	0.002*	0.004*	0.004*	0.053	-

Description: * $p < 0.05$ there is a significant difference

SPSS analysis showed that the antibacterial activity of the ethanol extract and the *n*-hexane fraction was not much different, while the ethanol extract and ethyl acetate fraction had quite different antibacterial activity. The

antibacterial activity of the ethyl acetate fraction and the positive control was not much different; this meant that the ethyl acetate fraction had good antibacterial activity against *Staphylococcus aureus* bacteria.

The results indicated that the activity as antibacterial of the ethanol extract against *Propionibacterium acne* belongs to the weak category, the *n*-hexane fraction belongs to the medium category, and the ethyl acetate fraction belongs to the strong category. Based on the results of the Shapiro-Wilk test, the value of $p > 0.05$, or the data is normally distributed, the test is continued with the homogeneity test, the result is $p < 0.05$, or the data is not homogeneous. The analysis was continued with a non-parametric test, namely Kruskal-Wallis, and continued with the Mann-Whitney test. The results of the Mann-Whitney test of the samples of *Garcinia forbesii* pericarp on the growth of *Propionibacterium acne* bacteria can be seen in Table 5.

Table 5. Mann-Whitney test results for the samples of *Garcinia forbesii* pericarp against *Propionibacterium acne* based on the diameter of the inhibition zone

Treatment group	Aquadest	Ethanol extract	<i>n</i> -hexane fraction	Ethyl acetate fraction	Chloramphenicol 1%
Aquadest	-	0.002*	0.002*	0.002*	0.002*
Ethanol extract	0.002*	-	0.055	0.004*	0.004*
<i>n</i>-hexane fraction	0.002*	0.055	-	0.004*	0.004*
Ethyl acetate fraction	0.002*	0.004*	0.004*	-	0.013*
Chloramphenicol 1%	0.002*	0.004*	0.004*	0.013*	-

Description: * $p < 0.05$ there is a significant difference

SPSS results showed that the activity as antibacterial of the ethanol extract and *n*-hexane fraction was not significantly different, while the antibacterial activity of the ethanol extract, *n*-hexane fraction, and ethyl acetate fraction to the positive control had quite a difference, so it can be interpreted that the resulting antibacterial activity of the ethanol extract, the *n*-hexane fraction and the ethyl acetate fraction were not as substantial as the antibacterial activity of the positive control which had been shown to inhibit bacterial growth because the positive control contained antibiotics.

4. Minimum Inhibitory Concentration Test Results (MIC)

MIC test results for the samples of *Garcinia forbesii* pericarp can be seen in Table 6. The results showed that the ethanol extract did not have MIC values for all bacteria, whereas for the *n*-hexane fraction, the MIC values were obtained for *Propionibacterium acne* at a concentration of 1.5%, and for *Staphylococcus aureus* bacteria, no MIC values were obtained, as well as for the ethyl acetate fraction obtained MIC values for *Propionibacterium acne* bacteria at a concentration of 1%, and MIC values were not obtained for *Staphylococcus aureus* bacteria. Based on the results acquired shows that the ethyl acetate fraction has the smallest MIC value compared to other samples; these results follow the outcomes of previous tests where the ethyl acetate fraction has the largest diameter of the inhibition zone compared to other samples.

Table 6. Minimum inhibitory concentration (MIC) of the samples of *Garcinia forbesii* pericarp

Sample	Concentration (%)	<i>Staphylococcus aureus</i>	<i>Propionibacterium acne</i>
Ethanol extract	1.5	+	+
	1.25	+	+
	1	+	+
	0.75	+	+
	0.5	+	+

<i>n</i>-hexane fraction	1.5	+	-*
	1.25	+	+
	1	+	+
	0.75	+	+
	0.5	+	+
Ethyl acetate fraction	1.5	+	-
	1.25	+	-
	1	+	-*
	0.75	+	+
	0.5	+	+
Positive control	1	-	-
Negative control		+	+

Information:

Sign (+) : there is bacterial growth

Sign (-) : no bacterial growth

Sign (-*): MIC value

5. Minimum Bactericidal Concentration Test Results (MBC)

The results of the MBC test for the ethanol extract, *n*-hexane fraction, and ethyl acetate fraction of *Garcinia forbesii* pericarp can be seen in Table 7. The results showed that the *n*-hexane fraction had a MBC value against *Propionibacterium acne* at a concentration of 1.5%, or the antibacterial activity of the *n*-hexane fraction against *Propionibacterium acne* was bactericidal. The ethyl acetate fraction obtained the MBC value for *Propionibacterium acne* at a concentration of 1.5% and was bacteriocidal, while for *Propionibacterium acne* at a concentration of 1%, it was bacteriostatic because, at that concentration, there was bacterial growth after incubation. This is because the higher the concentration, the more antibacterial compounds are contained in the sample, so the capability to inhibit bacterial growth is higher (Adila *et al*, 2013).

Table 7. Minimum bactericidal concentration (MBC) of ethanol extract, *n*-hexane fraction, and ethyl acetate fraction of *Garcinia forbesii* pericarp

Sample	Concentration (%)	<i>Staphylococcus aureus</i>	<i>Propionibacterium acne</i>
<i>n</i>-hexane fraction	1.5	Not tested	-*
	1.25	Not tested	Not tested
	1	Not tested	Not tested
ethyl acetate fraction	1.5	Not tested	-*
	1.25	Not tested	+
	1	Not tested	+

Information:

Sign (+) : there is bacterial growth

Sign (-) : no bacterial growth (MBC value)

Sign (-*): has no MIC value

CONCLUSION

The results of this study demonstrated that the ethyl acetate fraction of *Garcinia forbesii* pericarp was more effective than the ethanol extract and *n*-hexane fraction against *Propionibacterium acne* on the basis of zone of inhibition diameter, MIC and MBC. It has been shown to have strong antibacterial activity.

ACKNOWLEDGEMENT

Authors praise the blessings of God and thank to Faculty of Mathematics and Natural Sciences Universitas Lambung Mangkurat who has funded this research, and to everyone for their support and help.

REFERENCES

- Adila, rahmi, Nurmiati, N., Agustien, A., 2013. Uji Antimikroba Curcuma spp. Terhadap Pertumbuhan *Candida albicans*, *Staphylococcus aureus* dan *Escherichia coli*. *Jurnal Biologi Universitas Andalas* 2, 1–7.
- Alen, Y., Safitri, N., Dachriyanus, Manaf Ali, A., Ladjis, N.H., Sargent, M. V, 2008. Rubraxhantone Dari *Garcinia forbesii* KING. dan Bioaktivitasnya. *Jurnal Riset Kimia* 1, 192–201.
- Amalia, S., Wahdaningsih, S., Untari, E.K., 2014. Uji Aktivitas Antibakteri Fraksi *n*-Heksan Kulit Buah Naga Merah (*Hylocereus polyrhizus* Britton & Rose) Terhadap Bakteri *Staphylococcus aureus* ATCC 25923. *Jurnal Fitofarmaka Indonesia* 1.
- Anindya, D., 2012. Efek Ekstrak Kulit Buah Manggis (*Garcinia Mangostana* L.) Terhadap Pertumbuhan Bakteri *Shigella dysenteriae* dan *Escherichia coli*. UIN Syarif Hidayatullah, Jakarta.
- Asifa, U.S., 2014. Uji Aktivitas Antibakteri Fraksi *n*-Heksana Kulit Buah Manggis (*Garcinia mangostana* L.) Terhadap Pertumbuhan *Shigella Flexneri* Secara In Vitro. Universitas Tanjung Pura, Pontianak.
- Aziz, M.A., 2015. Uji Aktivitas Antibakteri Ekstrak Etanol kulit Buah Manggis (*Garcinia mangostana* l) Terhadap Bakteri *Escherichia coli* ATCC 11229 dan *Staphylococcus aureus* ATCC 6538 Secara In Vitro. Universitas Muhammadiyah Surakarta, Surakarta.
- Chikezie, I.O., 2017. Determination of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) using a novel dilution tube method. *Afr J Microbiol Res* 11, 977–980. <https://doi.org/10.5897/ajmr2017.8545>
- Dewi, A.R., 2018. Penetapan Kadar Fenol Dan Tanin Total Serta Uji Aktivitas Ekstrak Etanol Kulit Buah Mundar (*Garcinia forbesii* King.) Dengan Metode FRAP. Universitas Lambung Mangkurat, Banjarbaru.
- Firdausi, I., Retnowati, R., Sutrisno, 2015. Fraksinasi Ekstrak Metanol Daun Mangga Kasturi, (*Mangifera casturi* Koesterm) Dengan pelarut *n*-Butanol. *Kimia Student Journal* 1, 785–790.
- Kaharap, A.D., Mambo, C., Nangoy, E., 2016. Uji Efek Antibakteri Ekstrak Batang Akar Kuning (*Arcangelisia flava* Merr.) Terhadap Bakteri *Staphylococcus aureus* dan *Escherichia coli*. *Jurnal e-Biomedik (eBm)* 4.
- Karlina, C.Y., Ibrahim, M., Trimulyono, G., 2013. Aktivitas Antibakteri Ekstrak Herba Krokot (*Portulaca oleracea* L.) terhadap *Staphylococcus aureus* dan *Escherichia coli*. *Lentera Bio* 2, 87–93.
- Karthiga, P., Soranam, R., Annadurai, G., 2012. Alpha-mangostin, The Major Compound From *Garcinia mangostana* Linn. Responsible For Synthesis of Ag Nanoparticles: Its Characterization and Evaluation Studies. *Research Journal of Nanoscience and Nanotechnology* 2, 46–57.
- Lestari, Y., Ardiningsih, P., Nurlina, N., 2016. Aktivitas Antibakteri Gram Positif dan Negatif Dari Ekstrak Dan Fraksi Daun Nipah (*Nypa fruticans* Wurmb.) Asal Pesisir Sungai Kakap Kalimantan Barat. *Jurnal Kimia Khatulistiwa* 5, 1–8.

- Marsella, C., 2018. Penetapan Kadar Fenol dan Tanin Total serta Uji Aktivitas Antioksidan Fraksi Etil Asetat Kulit Buah Mundar (*Garcinia forbesii* King.) dengan Metode FRAP. Universitas Lambung Mangkurat, Banjarbaru.
- Miladiyah, I., Rachmawaty, F.J., 2017. Potency of Xanthone Derivatives as Antibacterial Agent Against Methicillin-Resistant *Staphylococcus Aureus* (MRSA). *Jurnal Kedokteran dan Kesehatan Indonesia* 8, 124–135.
- Mranani, S.A., 2015. Pemanfaatan Potensi Manggis Merah (*Garcinia forbesii*) Sebagai Pengawet Alami dan Minuman Fungsional. Institut Pertanian Bogor, Bogor.
- Noor, A., Ningsih, R.D., 2017. Mundar (*Garcinia forbesii*) Si Manggis Merah Sumber Daya Genetik Kalimantan Selatan. Banjarmasin.
- Noviardini, P.U., 2010. Uji Aktivitas Antibakteri Alfa Mangostin Kulit Buah Manggis (*Garcinia mangostana* L.) Terhadap Bakteri *Escherichia coli* Multiresisten Antibiotik dan Bakteri *Streptococcus* sp. Universitas Muhammadiyah Surakarta, Surakarta.
- Poeloengan, M., Praptiwi, P., 2010. Uji Aktivitas Antibakteri Ekstrak Kulit Buah Manggis (*Garcinia mangostana* Linn). *Media Litbang Kesehatan* XX, 65–69.
- Prasetyo, A.D., Sasongko, H., 2014. Aktivitas Antibakteri Ekstrak Etanol 70% Daun Kersen (*Muntingia calabura* L.) Terhadap Bakteri *Bacillus subtilis* dan *Shigella dysenteriae* Sebagai Materi Pembelajaran Biologi SMA Kelas X untuk Mencapai Kd 3.4 pada Kurikulum 2013. *Jurnal Penelitian Mahasiswa Pendidikan Biologi* 1, 98–102.
- Prihandani, S.S., Poeloengan, M., Noor, S.M., Andriani, 2015. Uji Daya Antibakteri Bawang Putih (*Allium sativum* L.) Terhadap Bakteri *Staphylococcus aureus*, *Escherichia coli*, *Salmonella typhimurium* dan *Pseudomonas aeruginosa* dalam Meningkatkan Keamanan Pangan. *Informatika Pertanian* 24, 53–58.
- Rachmawaty, D.U., 2016. Uji Aktivitas Antibakteri Ekstrak Etanol, Etil Asetat dan Petroleum Eter Rambut Jagung Manis (*Zea mays saccharata* Sturt) Terhadap Bakteri *Staphylococcus aureus* Dan *Escherichia Coli*. Universitas Islam Negeri Maulana Malik Ibrahim Malang, Malang.
- Ritthiwigrom, T., Laphookhieo, S., Pyne, S.G., 2013. Chemical constituents and biological activities of *Garcinia cowa* Roxb. *Maejo Int. J. Sci. Technol* 7, 212–231.
- Rizki, M.I., Nurlily, N., Fadlilaturrahmah, F., Ma'shumah, M., 2021. Skrining Fitokimia dan Penetapan Kadar Fenol Total Pada Ekstrak Daun Nangka (*artocarpus heterophyllus*), Cempedak (*Artocarpus integer*) dan Tarap (*Artocarpus odoratissimus*) Asal Desa Pengaron Kabupaten Banjar. *Jurnal Insan Farmasi Indonesia* 4, 95–102.
- Rostinawati, T., 2010. Aktivitas Antimikroba Ekstrak Herba Tespong (*Oenanthe javavica* D.C) Terhadap *Escherichia Coli*, *Staphylococcus Aureus* Dan *Candida albicans*. Jatinangor.
- Rosyad, P.G.Y., 2009. Formulasi Gel Obat Jerawat Minyak Atsiri Daun Jeruk Nipis (*Citrus aurantifolia*, Swingle) Dan Uji Daya Antibakteri (*Propionibacterium acne*) Secara In Vitro. Universitas Muhammadiyah Surakarta, Surakarta.
- Salni, S., Marisa, H., Mukti, R.W., 2011. Isolasi Senyawa Antibakteri Dari Daun Jengkol (*Pithecolobium lobatum* Benth) dan Penentuan Nilai KHM-nya. *Jurnal Penelitian Sains* 14, 38–41.
- Sambou, C.N., Wibowo, A.E., Taurhesia, S., 2017. Pengembangan Produk Sediaan Gel Kombinasi Ekstrak Daun Sirsak (*Annona muricata* L.) Dengan Ekstrak Rimpang Temulawak (*Curcuma xanthorrhiza* Roxb.) Sebagai Antibakteri Penyebab Jerawat (*Propionibacterium acne* dan *Staphylococcus epidermidis*). *PHARMACON Jurnal Ilmiah Farmasi - UNSRAT* 6, 255–265.
- Saputro, G.M.H., n.d. Uji Aktivitas Antibakteri Fraksi Etil Asetat Ekstrak Metanol Kulit Buah Manggis (*Garcinia mangostana* L.) Terhadap *Shigella flexneri*. Universitas Tanjung Pura, Pontianak.

- Sayuti, M., 2017. Pengaruh Perbedaan Metode Ekstraksi, Bagian dan Jenis Pelarut Terhadap Rendemen dan Aktivitas Antioksidan Bambu Laut (*Isis Hippuris*). *Technology Science and Engineering Journal* 1, 166–174.
- Sriyono, R.A.N., Andriani, I., n.d. Daya Antibakteri Ekstrak Etanol Kulit Manggis (*Garcinia Mangostana* Linn.) Terhadap Bakteri *Porphyromonas Gingivalis*.
- Wahyuddin, M., Pakadang, S.R., Aprilyani, A., 2017. Uji Aktivitas Anti Bakteri Ekstrak Etanol Daun Dewandaru (*Eugenia uniflora* L) Terhadap *Streptococcus pneumoniae* Dan *Shigella dysenteriae*. *Jurnal Farmasi UIN Alauddin Makassar* 5, 199–205.
- Weinstein, M.P., 2018. *Methods For Dilution Antimicrobial Susceptibility Tests For Bacteria That Grow Aerobically*, 11th ed. Clinical and Laboratory Standards Institute, Pennsylvania.
- World Health Organization, 2021. *Antimicrobial Resistance*. Geneva.

How to cite this Article: Izma, H., Rizki, M.I., Anwar, K., Anggraeni, D., Rahmatullah, S.W., Putra, A.M.P., Sandi, D.A.D., 2023. Antibacterial Activity of Ethanol Extract, n-Hexane and Ethyl Acetate Fraction of Mundar (*Garcinia forbesii*) Pericarp. *J. Pharm. Sci.* 6, 112-121.